

# EYEMOX-D

Eye Drops

Moxifloxacin HCl  
Dexamethasone Sodium Phosphate  
(Sterile Ophthalmic Solution)

## COMPOSITION

Each ml Contains:  
Moxifloxacin HCl equivalent to Moxifloxacin .....5mg & Dexamethasone Sodium Phosphate equivalent to Dexamethasone Phosphate .....1mg.  
(Vega Specs.)

## PRESERVATIVE

Banzalkonium Chloride.

## DOSAGE FORM

Ophthalmic solution

## DESCRIPTION

Eyemox-D is a combination of Moxifloxacin 0.5% and Dexamethasone Phosphate 0.1%. Moxifloxacin is a fourth generation Fluoroquinolone which controls infection by inhibiting the DNA Gyrase & Topoisomerase IV. Dexamethasone Phosphate a potent corticosteroid effectively controls the inflammation by inhibiting the release of inflammatory mediators.

## INDICATIONS

Eyemox-D is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists. The combination can also be used for post-operative inflammation and any other ocular inflammation associated with infection.

## DOSAGE

In the prevention of postoperative ocular infection and inflammation, instill 1 drop 3 times per day, in the eye to operate from 1 day before surgery until 15 days after surgery. In patients undergoing cataract surgery, day surgery, instill the medication immediately eye surgery or as prescribed.

In eye infections caused by susceptible microorganisms, instill 1 drop 3 times a day, 7 days or as medically indicated.

## CONTRAT-INDICATIONS

Eyemox-D is contraindicated in epithelial herpes simplex keratitis (Dendritic keratitis),

آئی موکس۔ ڈی  
موکسی فلوکسائین ایچ سی ایل  
ڈیکسامیٹھاسون سوڈیم فاسفیٹ

vaccinia, varicella, and in many other viral disease of the conjunctiva and cornea, mycobacterial infection of the eye and fungal diseases of ocular structures and in individuals hypersensitive to any of the components of the medication.

## WARNINGS & PRECAUTIONS

Prolonged use of steroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and field of visions, and posterior subcapsular cataract formation.

## ADVERSE EFFECTS

The most frequently reported drug-related undesirable effects seen with Moxifloxacin are conjunctival irritation, increased lacrimation, keratitis, papillary conjunctivitis and secondary infections due to steroid use.

## STORAGE

Store below 30°C.  
Protect from heat and sunlight.  
Keep out of the reach of children.  
On prescription only.  
Use the solution within four weeks after first opening the bottle.

## PRESENTATION

Eyemox-D eye drops is available in sterile plastic dropper bottler of 5 ml.

طریقہ استعمال:  
ایک قطرہ دن میں 3 مرتبہ متاثرہ آنکھ / آنکھوں میں ڈالیں  
یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔  
دوا کو 30 ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔  
دوا کو بچوں کی پہنچ سے دور رکھیں۔  
گرمی اور دھوپ سے بچائیں۔  
پہلی دفعہ بوتل کھولنے کے بعد دوا کو چار ہفتوں تک استعمال کیا جاسکتا ہے۔  
انتباہ:  
صرف مستند ڈاکٹر کے نسخہ پر استعمال کریں۔



Manufactured by:  
**Vega Pharmaceuticals (Pvt.) Ltd.**  
30 Km, Multan Road, Lahore, Pakistan.