

# VECHLOR

(CHLORAMPHENICOL)

Sterile Ophthalmic Solution

وی کلور  
کلورم فینی کول

## COMPOSITION:

Each ml contains:

Chloramphenicol ..... 5mg.

(B.P Specifications)

Hydroxypropyl Methylcellulose..... 5mg.

**Preservative:** Thiomersal

## ACTION:

Chloramphenicol is a broad spectrum antibiotic originally isolated from streptomyces venezuelae. It is primarily bacteriostatic and inhibits protein synthesis by interfering with the transfer of activated aminoacids from soluble RNA to ribosomes. Development of resistance to chloramphenicol can be regarded as minimal for Staphylococci and many other species of bacteria.

## INDICATIONS:

For the treatment of superficial ocular infection involving the conjunctiva and/or cornea caused by chloramphenicol-susceptible organisms. Bacteriological studies should be performed to determine the causative organisms and their sensitivity to chloramphenicol.

## CONTRAINDICATIONS:

Contraindicated in those individuals who have shown hypersensitivity to any of its components.

## WARNINGS:

Long term or frequent intermittent use of topical chloramphenicol should be avoided because of the possibility of causing hypersensitivity reactions including bone-marrow hypoplasia which may lead to aplastic anemia.

## PRECAUTIONS:

Long term use of antibiotic may occasionally result in overgrowth of non-

susceptible organisms including fungi. If new infections appear during treatment, the drug should be discontinued and appropriate measures should be taken in all except very superficial infections. The topical use of chloramphenicol should be supplemented with appropriate systemic medication.

## ADVERSE REACTIONS:

Blood dyscrasias may be associated with the systemic use of chloramphenicol. One case of bone-marrow hypoplasia following the long term (23 months) topical use of chloramphenicol ophthalmic solution has been reported.

## ADMINISTRATION:

1 to 2 drops 4 to 6 times a day for the first 72 hours, depending upon the severity of the condition. Since the action of the drug is primarily bacteriostatic, therapy should be continued for 48 hours after an apparent cure has been attained.

## HOW SUPPLIED:

As 10 ml sterile ophthalmic solution in plastic dropper bottles.

## INSTRUCTIONS:

Store below 30°C.

Protect from heat and sunlight.

Keep out of the reach of children.

Use only on medical advice.

For Ophthalmic use only.

Use within four weeks after first opening.

خوراک: ڈاکٹر کی ہدایات کی مطابق استعمال کریں۔

ہدایات: دوا کو دھوپ اور گرمی سے بچائیں۔

۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔

تمام ادویات بچوں کی پہنچ سے دور رکھیں۔

پہلی دفعہ بوتل کھولنے کے بعد دوا کو چار ہفتوں تک استعمال کیا جاسکتا ہے۔



Manufactured by:

**Vega Pharmaceuticals (Pvt.) Ltd.**

30 Km, Multan Road, Lahore, Pakistan.